



Sagebrush Ecosystem Program

Findings and Improvement Recommendations

April 5th, 2017

Sagebrush Ecosystem Technical Team (SETT)

Kelly McGowan, Program Manager

Katie Andrle, NDOW

Dan Huser, NDF

Sara McBee, NDSL

Vacant, NDA



Research – Raven Abundance and Occupancy

- Probability of occupancy (nesting) on anthropogenic structures
 - 80% (53% Transmission Lines; 17% cell towers; 4% nesting platforms; Coates et al. 2014)
 - 90% within 400m of survey locations at oil development, cities, and roads (Bui et al. 2010)
- Howe et al. (2014) found a 31% decrease in odds of nesting ravens for every 1km increase in distance from a transmission line
- Increased raven nesting occupancy by 80% following construction of a 500kV transmission line over 9 years (Steenhof et al. 1993)



Research – Raven Abundance and Occupancy

- Coates et al. (2014) determined the probability of raven occurrence in relation to transmission lines was most significant within 2.2 km of a line, but effect was observed out to 27 km
- Bui et al. (2010) found that raven density was highest within 3km of cities, urban areas, and associated infrastructure and dropped sharply beyond this distance
- Gibson et al. (in Review) linked impacts on sage-grouse behavior and demographics to indirect effects of raven colonization of the Falcon-Gondor (FG) line out to 10km



Research – Raven Nest Depredation

- Ravens have been identified as the primary nest predator of sage-grouse in multiple studies in Nevada and Idaho (Coates et al. 2008, Coates and Delehanty 2008, Coates and Delehanty 2010, Lockyer et al. 2013)
- Sage-grouse survival in Wyoming was more affected by occupancy (e.g. nesting, territorial ravens) than raven density (non-territorial, nomadic ravens) and were responsible for the majority of nest depredations (Bui et al. 2010)



Research – Sage-grouse Avoidance of Transmission Lines

- Sage-grouse showed avoidance within 600m of transmission lines (138kV) in Idaho based on telemetry studies (Gillian et al. 2013), 500m based on pellet counts (Braun 1998, Hanser et al. 2011)
- Hansen et al. (2016) demonstrated transmission line presence negatively influenced sage-grouse winter habitat use in Utah
 - Did not find a difference in use/avoidance between pre- and post-construction of adjacent 345kV line
- Pruett et al. (2008) also found avoidance of transmission lines by greater and lesser prairie chickens in Oklahoma
- Resistance models parameterized by expert opinion that were developed in Washington state predicted that powerlines would significantly affect sage-grouse movement, gene flow, and lek activity to distances beyond 500m (Shirk et al. 2015), which can lead to population level impacts



Research – Sage-grouse Demographics and Transmission Lines

- Gibson et al. (In Review)
 - Nests within 8km of FG line had reduced nest success
 - Pre-fledgling chick survival was lower closer to the line
 - Population growth rate, measure by lek attendance, declined by 3% within 10km
 - Nest site selection was reduced within 10km of the line
- Wisdom et al. (2011) reported the mean distance of historical sage-grouse locations to transmission lines was 6km in extirpated range compared to 15km in currently occupied range
- Dinkins et al. (2014) showed an increase in female mortality with increased density of powerlines
- LeBeau et al. (2014) reported nest survival was not influenced by distance to transmission lines
 - The relationship was not substantial due to very large 90% confidence intervals
 - Habitat quality also influenced and confounded results – nest survival was negatively associated with lower variation in shrub height



Examples of Applied Science in Other States

- FWS and BLM representatives convened to assess the indirect effects of transmission lines and associated infrastructure, produced Whitepaper in 2015
 - “Assessing indirect effects of transmission lines on greater sage-grouse for the Gateway West Interstate Transmission Line Project”
- Described three indirect impact zones
 - Avoidance (600m impact zone) – recommended 75-95% habitat services reduction
 - Increased Avian Predator Presence and Predation (600-1,200m impact zone) – recommended a 20-50% habitat services reduction
 - Decreased Productivity and Survival (1,200-5,000m impact zone) – recommended a 5-40% habitat services reduction



Examples of Applied Science in Other States

- Selection of six members for a Technical Advisory Group to develop a science based approach to quantify indirect effects of transmission lines for two projects: Energy Gateway South and TransWest in WY, CO, UT. FEIS available December 2016
- Revised conclusions in 2015 Whitepaper with latest available research
 - Avoidance (600m impact zone)
 - Decreased Population Growth (0-10,000m impact zone)
- Determined habitat services lost declined linearly to 10km with 75% reduction at the line



Improvement Recommendation

DISTURBANCE TYPE	Disturbance Subtype	WEIGHT (%)	DISTANCE (Kilometers)
Towers (cell, etc.)	NA	75%	6 km
Power Lines	Transmission	75%	6 km
Power Lines	Distribution	25%	6 km

- Transmission Lines
 - High Voltage
 - Steel Lattice, H frame, monopole with cross members
- Distribution Lines
 - Monopole with no cross members, supporting arms, etc, or of a construction that would not support nesting that can be documented
- Primary reasoning for sub-diving relates to ability of ravens to nest and occupy lines, which would include minimization measures (e.g. effective perch deterrents)



Improvement Recommendation

DISTURBANCE TYPE	Disturbance Subtype	WEIGHT (%)	DISTANCE (Kilometers)
Towers (cell, etc.)	NA	75%	6 km
Power Lines	Transmission	75%	6 km
Power Lines	Distribution	25%	6 km


Recommendation	Rationale
75% Weight	<ol style="list-style-type: none"> 1) Avoidance – 600m avoidance zone received 75% Habitat Services Reduction (FWS/BLM 2015, BLM 2016) 2) Raven Occupancy – 80% (Coates et al. 2014); 90% (Bui et al. 2010)
25% Weight	<ol style="list-style-type: none"> 1) Probability of Raven Occurrence – 30% (Coates et al. 2014) 2) Effective minimization measures to significantly reduce raven occupancy (Slater and Smith and Smith 2010; Dwyer and Doloughan 2014; Dwyer and Leiker 2012)
6 km	<ol style="list-style-type: none"> 1) Reduced demographic rates Gibson et al. (In Review) <ul style="list-style-type: none"> • Reduced Lambda (population growth) by 3% - 10km • Reduced nest survival - 8km • Reduced pre-fledgling chick survival 2) Distance – Highest probability of impact <ul style="list-style-type: none"> • Coates et al. (2014) – 2.2km, extend to 27 • Bui et al. (2010) – within 3km



Improvement Recommendation


DISTURBANCE TYPE	Disturbance Subtype	WEIGHT (%)	DISTANCE (Kilometers)
Towers (cell, etc.)	NA	75%	6 km
Power Lines	Transmission	75%	6 km
Power Lines	Distribution	25%	6 km

- Change from 8 to 6km to reflect the most significant impacts to sage-grouse (within 3km)
- The impact measured by the HQT beyond 6km (when using 8km distance) is minimal (~1%)





Improvement Recommendation



Distance (m)	Weight
0	75%
800	57%
1600	40%
2400	27%
3200	17%
4000	8%
5000	2%
6000	0%